

XIAFLEX[®]
collagenase clostridium histolyticum injection
0.9mg

//// THE XIAFLEX COPAY
ASSISTANCE PROGRAM

~94%

OF ELIGIBLE
PATIENTS SHOULD
PAY \$0 OUT OF
POCKET FOR
XIAFLEX^{*,†}

What you pay for XIAFLEX depends on your health benefits or insurance coverage. Look inside to see if you qualify.

*Most eligible patients with commercial insurance plans should pay a \$0 copay for XIAFLEX. Predictions based on historical analysis of claims filed September 2022 through August 2023.

†See inside for Restrictions, Terms and Conditions.

Please see Important Safety Information on pages 3-4, and click for [full Prescribing Information](#), including **BOXED WARNING and [Medication Guide](#).**

IF ELIGIBLE, YOU MAY PAY \$0 OUT OF POCKET FOR XIAFLEX^{*,†}

You are eligible for assistance under the XIAFLEX Copay Assistance Program if:

- You are receiving or have received XIAFLEX for the approved indication and in a manner consistent with the instructions for administration of XIAFLEX
- You are uninsured or have insurance that is not provided by Medicare, Medicare Prescription Drug Benefit plans, Medicare Advantage, Veterans Affairs (VA), Medicaid, or similar federal or state programs, and this program is not otherwise prohibited by law
- You are 18 years of age or older
- You have paid or are obligated to pay out-of-pocket costs for a prescription of XIAFLEX

No other purchase is necessary to receive this offer.

No physical savings card is required for this copay program.

^{*}Most eligible patients with commercial insurance plans should pay a \$0 copay for XIAFLEX. Predictions based on historical analysis of claims filed September 2022 through August 2023.

[†]Restrictions, Terms and Conditions

- By accepting this offer, you agree to report the value received under this offer to any health insurer or other third party paying for any part of the XIAFLEX prescription if you are required to do so by benefit terms, contract, or law.
- This offer is not valid for prescriptions reimbursed in whole or in part by Medicare, Medicare Prescription Drug Benefit plans, Medicare Advantage, VA, Medicaid, or similar federal or state programs, or where otherwise prohibited by law.
- By accepting this offer, you agree that Endo Pharmaceuticals Inc. or those working on its behalf may contact your doctor to verify information about treatment that is relevant to verifying your eligibility for this offer.
- This offer is only valid for doses of XIAFLEX administered in the U.S.
- This offer is valid for the out-of-pocket cost for the dose of XIAFLEX only. Offer is not valid for any other products or other out-of-pocket costs (for example, office visit charges, office visit copays, or injection/administration costs), even if those costs are associated with the administration of a dose of XIAFLEX.
- This offer is valid only if you have not used this program within the last 30 days.
- The selling, purchasing, trading, or counterfeiting of this offer is prohibited.
- Endo Pharmaceuticals Inc. reserves the right to rescind, revoke, or amend this offer without notice.
- By participating, you understand and agree to comply with the terms and conditions of this offer as set forth above.



To find out if you're eligible, scan
the QR code or call 1-800-743-2382

WHAT IS XIAFLEX® (collagenase clostridium histolyticum)?

XIAFLEX is a prescription medicine used to treat adult men with Peyronie's disease who have a "plaque" that can be felt and a curve in their penis greater than 30 degrees when treatment is started.

It is not known if XIAFLEX is safe and effective in children under the age of 18.

IMPORTANT SAFETY INFORMATION

Do not receive XIAFLEX if:

- the Peyronie's plaque to be treated involves the "tube" that your urine passes through (urethra).
- you are allergic to collagenase clostridium histolyticum or any of the ingredients in XIAFLEX, or to any other collagenase product. See the end of the Medication Guide for a complete list of ingredients in XIAFLEX.

XIAFLEX can cause serious side effects, including:

1. **Penile fracture (corporal rupture) or other serious injury to the penis.** Receiving an injection of XIAFLEX may cause damage to the tubes in your penis called the corpora. After treatment with XIAFLEX, one of these tubes may break during an erection. This is called a corporal rupture or penile fracture. This could require surgery to fix the damaged area. Damage to your penis might not get better after a corporal rupture.
 - After treatment with XIAFLEX, blood vessels in your penis may also break, causing blood to collect under the skin (hematoma). This could require a procedure to drain the blood from under the skin. If a hematoma appears, skin and soft tissue necrosis (death of skin cells) may develop in that area, which could require surgery.

Symptoms of corporal rupture or other serious injury to your penis may include:

- a popping sound or sensation in an erect penis
- sudden loss of the ability to maintain an erection
- pain in your penis
- purple bruising and swelling of your penis
- difficulty urinating or blood in the urine

Call your healthcare provider right away if you have any of the symptoms of corporal rupture or serious injury to the penis listed above.

Do not have sex or any other sexual activity between the first and second injections of a treatment cycle.

Do not have sex or have any other sexual activity for at least 4 weeks after the second injection of a treatment cycle with XIAFLEX and after any pain and swelling has gone away.

XIAFLEX for the treatment of Peyronie's disease is only available through a restricted program called the XIAFLEX Risk Evaluation and Mitigation Strategy (REMS) Program.

2. **Hypersensitivity reactions, including anaphylaxis.** Severe allergic reactions can happen in people who receive XIAFLEX, because it contains foreign proteins.

Call your healthcare provider right away if you have any of these symptoms of an allergic reaction after an injection of XIAFLEX:

- hives
- chest pain
- swollen face
- low blood pressure
- breathing trouble
- dizziness or fainting

3. **Back pain reactions.** After receiving an injection of XIAFLEX for Peyronie's disease, you may suddenly feel back pain, including severe lower back pain moving to your legs, feet, chest and arms. The back pain may also include spasms and make it hard to walk. These symptoms usually go away in 15 minutes or less, but may last longer.

Tell your healthcare provider right away if you have sudden back pain, chest pain, or hard time walking after an injection.

(continued on next page)

IMPORTANT SAFETY INFORMATION (CONT.)

4. Fainting. Fainting (passing out) or near fainting can happen in men who receive XIAFLEX®, especially if they have severe penile pain.

If you have dizziness or feel faint after receiving XIAFLEX, lie down until the symptoms go away.

Before receiving XIAFLEX, tell your healthcare provider if you have had an allergic reaction to a previous XIAFLEX injection, have a bleeding problem, received XIAFLEX for another condition, or any other medical conditions. Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Using XIAFLEX with certain other medicines can cause serious side effects. Especially tell your healthcare provider if you take medicines to thin your blood (anticoagulants). If you are told to stop taking a blood thinner before your XIAFLEX injection, your healthcare provider should tell you when to restart the blood thinner. Ask your healthcare provider or pharmacist for a list of these medicines, if you are not sure.

What should I avoid while receiving XIAFLEX?

Avoid situations that may cause you to strain your stomach (abdominal) muscles, such as straining during bowel movements.

Do not use a vacuum erection device during your treatment with XIAFLEX.

XIAFLEX can cause serious side effects, including increased chance of bleeding. Bleeding or bruising at the injection site can happen in people who receive XIAFLEX. Talk to your healthcare provider if you have a problem with your blood clotting. XIAFLEX may not be right for you.

The most common side effects with XIAFLEX for the treatment of Peyronie's disease include:

- a small collection of blood under the skin at the injection site (hematoma)
- swelling at the injection site or along your penis
- pain or tenderness at the injection site, along your penis and above your penis
- penis bruising
- itching of your penis or scrotum (genitals)
- painful erection
- erection problems (erectile dysfunction)
- changes in the color of the skin of your penis
- blisters at the injection site
- pain with sex
- a lump at the injection site (nodule)

Tell your healthcare provider if you have any side effect that bothers you or does not go away.

These are not all of the possible side effects with XIAFLEX. For more information, ask your healthcare provider or pharmacist. You may report side effects to FDA at 1-800-FDA-1088.

Click for [full Prescribing Information](#), including [BOXED WARNING](#) and [Medication Guide](#).